

Scheldeman, L. et al. (2019) Total mismatch in diffusion negative patients in the WAKE-UP trial. *International Journal of Stroke*, 14(7), NP20-NP22. (doi: [10.1177/1747493019875238](https://doi.org/10.1177/1747493019875238)).


This is the author's final accepted version.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/190027/>

Deposited on: 11 July 2019

Enlighten – Research publications by members of the University of Glasgow  
<http://eprints.gla.ac.uk>



### **Total Mismatch in Diffusion Negative Patients in the WAKE-UP Trial**

Journal:	<i>International Journal of Stroke</i>
Manuscript ID	IJS-06-19-7225
Manuscript Type:	Letter to the Editor
Date Submitted by the Author:	24-Jun-2019
Complete List of Authors:	Scheldeman, Lauranne; KU Leuven University Hospitals Leuven

	<p>Gasthuisberg Campus, Neurology Wouters, Anke; UZLeuven, Neurology Dupont, Patrick; KULeuven, Laboratory for Epilepsy Research, Laboratory for Cognitive Neurology; University Hospitals Leuven, Medical Imaging Research Center Cheng, Bastian; University Medical Center Hamburg-Eppendorf, Neurology Ebinger, Martin; Medical Park Humboldtmühle, Neurology Endres, Matthias; Charité Universitätsmedizin Berlin Campus Charite Mitte, Centrum für Schlangenanfallforschung Berlin (CSB) Fiebach, Jochen; Charité University Hospital, Center of Stroke Research Berlin Gerloff, Christian; University Medical Center Hamburg-Eppendorf, Neurology Muir, Keith; University of Glasgow, Centre for Stroke and Brain Imaging Research, Institute of Neuroscience and Psychology; Southern General Hospital, Institute of Neurological Sciences Nighoghossian, Norbert; Université de Lyon 1, Stroke Unit, Hôpital Neurologique Pierre Wertheimer, Neurology Pedraza, Salvador; Institut de Diagnostic per la Image (IDI), Department of Radiology Simonsen, Claus; Aarhus University, Neurology Boutitie, Florent; Hospices Civils de Lyon, Service de Biostatistique Thomalla, Götz; Department of Neurology, University Medical Center Hamburg-Eppendorf (UKE) Lemmens, Robin; KULeuven, Experimental Neurology; University Hospitals Leuven, Neurology</p>
Keywords:	<p>Total mismatch, Magnetic Resonance Imaging, Ischemic Stroke, Diffusion Weighted Imaging, Perfusion Weighted Imaging, Fluid Attenuated Inversion Recovery</p>

**Total Mismatch in Diffusion Negative Patients in the WAKE-UP Trial**

Lauranne Scheldeman MD<sup>1-3</sup>, Anke Wouters MD PhD<sup>1-3</sup>, Patrick Dupont PhD<sup>4</sup>, Bastian Cheng MD<sup>5</sup>, Martin Ebinger MD<sup>6,7</sup>, Matthias Endres MD<sup>6,8</sup>, Jochen B. Fiebach MD<sup>6</sup>, Christian Gerloff MD<sup>5</sup>, Keith W. Muir MD<sup>9</sup>, Norbert Nighoghossian MD<sup>10</sup>, Salvador Pedraza MD, PhD<sup>11</sup>, Claus Z. Simonsen MD, PhD<sup>12</sup>, Florent Boutitie PhD<sup>13</sup>, Götz Thomalla MD<sup>5</sup>, Robin Lemmens MD PhD<sup>1-3</sup>

<sup>1</sup>Department of Neurology, University Hospitals Leuven, Herestraat 49, 3000 Leuven, Belgium

<sup>2</sup>KU Leuven – University of Leuven, Department of Neurosciences, Experimental Neurology, Oude Markt 13, bus 5005, 3000 Leuven, Belgium

<sup>3</sup>VIB, Center for Brain & Disease Research, Laboratory of Neurobiology, Campus Gasthuisberg, Herestraat 49, bus 602, 3000 Leuven, Belgium

<sup>4</sup>KU Leuven – University of Leuven, Department of Neurosciences, Laboratory for cognitive neurology, Leuven, Belgium

<sup>5</sup>Klinik und Poliklinik für Neurologie, Kopf- und Neurozentrum, University Medical Center Hamburg-Eppendorf, Martinistr. 52, 20246 Hamburg, Germany

<sup>6</sup>Centrum für Schlaganfallforschung Berlin (CSB), Charité - Universitätsmedizin Berlin, Campus Mitte, Charitéplatz 1, 10117 Berlin, Germany

<sup>7</sup>Neurologie der Rehaklinik Medical Park Humboldtmühle, An der Mühle 2-9, 13507 Berlin, Germany

<sup>8</sup>Klinik und Hochschulambulanz für Neurologie, Charité-Universitätsmedizin Berlin, Campus Mitte, Charitéplatz 1, 10117 Berlin, Germany

<sup>9</sup>Institute of Neuroscience & Psychology, University of Glasgow, University Avenue, Glasgow G12 8QQ, UK

<sup>10</sup>Department of Stroke Medicine, Université Claude Bernard Lyon 1, CREATIS CNRS UMR 5220-INSERM U1206, INSA-Lyon; Hospices Civils de Lyon, Lyon, France

<sup>11</sup>Department of Radiology, Institut de Diagnostic per la Image (IDI), Hospital Dr Josep Trueta, Institut d'Investigació Biomèdica de Girona (IDIBGI), Parc Hospitalari Martí i Julià de Salt - Edifici M2, 17190 Salt, Girona, Spain

<sup>12</sup>Department of Neurology, Aarhus University Hospital, 8200 Aarhus N, Denmark

<sup>13</sup>Hospices Civils de Lyon, Service de Biostatistique, F-69003 Lyon, France; Université Lyon 1, F-69100 Villeurbanne, France; CNRS, UMR 5558, Laboratoire de Biométrie et Biologie Evolutive, Equipe Biostatistique-Santé, F-69100 Villeurbanne, France

### **Corresponding author**

Lauranne Scheldeman

Neurology Department

University Hospitals Leuven

Herestraat 49

B-3000

Belgium

Email: [Lauranne.scheldeman@uzleuven.be](mailto:Lauranne.scheldeman@uzleuven.be)

Phone: +32163449054

Fax: +3216344285

**Key words:** Ischemic Stroke, Magnetic resonance imaging, Total mismatch

### **Word count:**

- Total: 1051
- Manuscript (excl. title page, acknowledgments, disclosures, funding and references): 438

**Figures:** 1

Dear Editor,

The WAKE-UP trial screened patients in the unknown time window after stroke onset for the presence of a DWI-FLAIR mismatch <sup>1</sup>. Since perfusion-weighted imaging (PWI) was not mandatory, the pattern of a total mismatch, i.e. a negative DWI together with a positive PWI lesion, remained undetected in the primary analysis <sup>2-6</sup>.

In a post-hoc analysis, we investigated the incidence of the total mismatch profile in patients enrolled in the WAKE-UP trial. For patients with available PWI, we calculated DWI and PWI lesion volumes with RAPID (iSchemaView, Menlo Park, CA). Of 1,362 patients enrolled, PWI was available in 343 (25%). Of these, only 3 (0.9%) had a total mismatch caused by a M2 occlusion of the middle cerebral artery (Figure 1).

**Figure 1. DWI and PWI analysis in total mismatch patients**

*[Insert figure 1.]*

DWI (A), PWI lesion with a time to maximum of the residue function (Tmax) >6s (B), PWI lesions with different Tmax value (C) generated by RAPID software (iSchemaView) and time-of-flight magnetic resonance angiography (D). For patient 1, 2 and 3, time from symptom recognition to imaging was 80, 269 and 90 min, time from last seen well to imaging was 410, 899 and 510 min and PWI lesion volume was 70, 76 and 32 ml respectively. All patients had an M2 occlusion of the left middle cerebral artery.

Less than one percent of patients had a total mismatch. The absence of a DWI lesion in patients presenting with stroke-like symptoms should prompt the clinician to consider stroke mimics. Additional PWI can guide physicians to determine a vascular etiology for the neurological symptoms, which can be confirmed by identifying a vessel occlusion on magnetic resonance

angiography. Apart from stroke mimics, early imaging after stroke onset, stroke in the posterior circulation and stroke caused by small vessel disease can all lead to DWI negativity.<sup>7</sup> None of these were present in the 3 total mismatch patients. The confirmation of a vessel occlusion excluded early (partial) reperfusion as a possible explanation for DWI negativity.

Since randomization in the WAKE-UP trial was based on the presence of the DWI-FLAIR mismatch<sup>1</sup>, these patients were not randomized and assigned to a particular treatment arm. Given the infinite PWI-DWI mismatch ratio, they might have been ideal candidates for reperfusion treatment. The natural course and benefit from treatment in these patients is not well known. Withholding reperfusion treatment might therefore be hazardous as long as the fate of the hypoperfused tissue remains uncertain.<sup>2-6</sup>

When evaluating patients for the DWI-FLAIR mismatch, PWI can provide additional information to ensure the most comprehensive patient selection, although the total mismatch pattern may be uncommon.

## Acknowledgments

None

## Conflicts-of-Interest/disclosures

Claus Z. Simonsen: Funded by a grant from Novo Nordisk Foundation.

Götz Thomalla: Consulting fees from Acandis, grant support and lecture fees from Bayer, lecture fees from Boehringer Ingelheim, Bristol-Myers Squibb/Pfizer, and Daiichi Sankyo, and consulting fees and lecture fees from Stryker.


## Sources of Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The WAKE-UP trial received funding from the European Union Seventh Framework Program [FP7/2007-2013] under grant agreement No. 278276.

## References

1. Thomalla G, Simonsen CZ, Boutitie F, et al. MRI-Guided Thrombolysis for Stroke with Unknown Time of Onset. *N Engl J Med* 2018; 379: 611-622. 2018/05/17. DOI: 10.1056/NEJMoa1804355.
2. Cho TH, Hermier M, Alawneh JA, et al. Total mismatch: negative diffusion-weighted imaging but extensive perfusion defect in acute stroke. *Stroke* 2009; 40: 3400-3402. 2009/08/22. DOI: 10.1161/STROKEAHA.109.563064.
3. Neeb L, Geisler F, Wendt M, et al. Thrombolytic therapy in total mismatch with severe stroke after acute MCA-occlusion and negative DWI. *Clin Neurol Neurosurg* 2013; 115: 802-804. 2012/08/09. DOI: 10.1016/j.clineuro.2012.07.018.
4. Kim JT, Park MS, Kim MK, et al. Minor stroke with total mismatch after acute MCA occlusion. *J Neuroimaging* 2011; 21: 399-402. 2011/01/13. DOI: 10.1111/j.1552-6569.2010.00564.x.
5. Hotter B, Ostwaldt AC, Levichev-Connolly A, et al. Natural course of total mismatch and predictors for tissue infarction. *Neurology* 2015; 85: 770-775. 2015/08/02. DOI: 10.1212/WNL.0000000000001889.
6. Fustier A, Naggara O, Tisserand M, et al. Total mismatch in anterior circulation stroke patients before thrombolysis. *J Neuroradiol* 2013; 40: 158-163. 2013/06/29. DOI: 10.1016/j.neurad.2013.03.001.
7. Edlow BL, Hurwitz S and Edlow JA. Diagnosis of DWI-negative acute ischemic stroke: A meta-analysis. *Neurology* 2017; 89: 256-262. 2017/06/16. DOI: 10.1212/WNL.0000000000004120.





DWI (A), PWI lesion with a time to maximum of the residue function (Tmax) >6s (B), PWI lesions with different Tmax value (C) generated by RAPID software (iSchemaView) and time-of-flight magnetic resonance angiography (D). For patient 1, 2 and 3, time from symptom recognition to imaging was 80, 269 and 90 min, time from last seen well to imaging was 410, 899 and 510 min and PWI lesion volume was 70, 76 and 32 ml respectively. All patients had an M2 occlusion of the left middle cerebral artery.